

A Holstein-Primakoff and a Dyson realization for the quantum algebra $U_q[sl(n+1)]$

Tchavdar D. Palev*

Department of Physics and Astronomy, University of Rochester, Rochester, New York 14627

Abstract. The known Holstein-Primakoff and Dyson realizations of the Lie algebra $sl(n+1)$, $n = 1, 2, \dots$ in terms of Bose operators (Okubo S 1975 *J. Math. Phys.* **16** 528) are generalized to the class of the quantum algebras $U_q[sl(n+1)]$ for any n . It is shown how the elements of $U_q[sl(n+1)]$ can be expressed via n pairs of Bose creation and annihilation operators.

In the present note we write down an analogue of the Dyson (D) realization and of the Holstein-Primakoff (HP) realization for the quantum algebra $U_q[sl(n+1)]$. Initially the HP and the D realizations were given for $sl(2)$ [1, 2]. The generalization for $gl(n+1)$ is due to Okubo [3]. In the realization of Okubo the elements of $gl(n+1)$ are expressed as functions of n pairs of Bose creation and annihilation operators (CAOs), namely operators $a_1^\pm, a_2^\pm, \dots, a_n^\pm$, which satisfy the known commutation relations

$$[a_i^-, a_j^+] = \delta_{ij}, \quad [a_i^+, a_j^+] = [a_i^-, a_j^-] = 0, \quad i, j = 1, \dots, n. \quad (1)$$

This realization is “more economical” than the known Jordan-Schwinger realization, which expresses $gl(n+1)$ via $n+1$ pairs of Bose CAOs.

The motivation in the present work stems from the various applications of the Holstein-Primakoff and of the Dyson realizations in theoretical physics. Beginning with [1] and [2] the HP and D realizations were constantly used in condensed matter physics. Some other early applications can be found in the book of Kittel [4] (more recent results are contained in [5]). For applications in nuclear physics see [6, 7] and the references therein, but there are, certainly, several other publications. In view of the importance of the

* Permanent address: Institute for Nuclear Research and Nuclear Energy, 1784 Sofia, Bulgaria; e-mail: tpalev@inrne.acad.bg

quantum algebras for physics, one could expect that the generalization of the Dyson and of the Holstein-Primakoff realizations to the case of quantum algebras may be of interest too. This is in fact the case for the only known so far q -analogs of the HP realization, namely those of $U_q[sl(2)]$ and $U_q[sl(3)]$ [8-15].

To begin with we recall the definition of $U_q[sl(n+1)]$ in the sense of Drinfeld [16]. Let $\mathbf{C}[[h]]$ be the complex algebra of the formal power series in the indeterminate h , $q = e^{h/2} \in \mathbf{C}[[h]]$. Then $U_q[sl(n+1)]$ is a Hopf algebra, which is a topologically free $\mathbf{C}[[h]]$ module (complete in the h -adic topology), with generators $\{h_i, e_i, f_i\}_{i=1, \dots, n}$ and

1. Cartan relations

$$[h_i, e_j] = (2\delta_{ij} - \delta_{i,j-1} - \delta_{i-1,j})e_j, \quad (2a)$$

$$[h_i, f_j] = -(2\delta_{ij} - \delta_{i,j-1} - \delta_{i-1,j})f_j, \quad (2b)$$

$$[e_i, f_j] = \delta_{ij} \frac{q^{h_i} - \bar{q}^{h_i}}{q - \bar{q}}. \quad (2c)$$

2. Serre relations

$$[e_i, e_j] = 0, \quad [f_i, f_j] = 0, \quad |i - j| \neq 1, \quad (3a)$$

$$[e_i, [e_i, e_{i\pm 1}]_{\bar{q}}]_q = [e_i, [e_i, e_{i\pm 1}]_q]_{\bar{q}} = 0, \quad (3b)$$

$$[f_i, [f_i, f_{i\pm 1}]_{\bar{q}}]_q = [f_i, [f_i, f_{i\pm 1}]_q]_{\bar{q}} = 0. \quad (3c)$$

Above and throughout $[a, b] = ab - ba$, $[a, b]_x = ab - xba$, $\bar{q} = q^{-1}$. We do not write the other Hopf algebra maps (Δ, ε, S) , since we will not use them. They are certainly also a part of the definition.

The Dyson and the Holstein-Primakoff realizations are different embeddings of $U_q[sl(n+1)]$ into the Weyl algebra $W(n)$. We define the latter as a topologically free $\mathbf{C}[[h]]$ module and an associative unital algebra with generators $a_1^\pm, \dots, a_n^\pm$ and relations (1).

Remark. In the physical applications it is often more convenient to consider h and q as complex numbers, $h, q \in \mathbf{C}$. Then all our considerations remain true provided q is not a root of 1. The replacement of $q \in \mathbf{C}[[h]]$ with a number corresponds to a factorization of $U_q[sl(n+1)]$ and $W(n)$ with respect to the ideals generated by the relation $q = \text{number}$. The factor-algebras $U_q[sl(n+1)]$ and $W(n)$ are complex associative algebras. However the completion in the h -adic topology has left a relevant trace: after the factorization the elements of $U_q[sl(n+1)]$ and of $W(n)$ are not only polynomials of their generators. In particular the functions of the CAOs, which appear in the D and in the HP realizations (see (4) and (10) bellow) are well defined as elements from $W(n)$.

Now we are ready to state our main results. Let $[x] = \frac{q^x - \bar{q}^x}{q - \bar{q}}$, $N_i = a_i^+ a_i^-$ and $N = N_1 + \dots + N_n$.

Proposition 1 (Dyson realization). The linear map $\varphi : U_q[sl(n+1)] \rightarrow W(n)$, defined on the generators as

$$\varphi(h_1) = p - N - N_1, \quad \varphi(h_i) = N_{i-1} - N_i, \quad i = 2, 3, \dots, n, \quad (4a)$$

$$\varphi(e_1) = \frac{[N_1 + 1]}{N_1 + 1} [p - N] b_1^-, \quad \varphi(e_i) = \frac{[N_i + 1]}{N_i + 1} b_i^- b_{i-1}^+, \quad i = 2, \dots, n, \quad (4b)$$

$$\varphi(f_1) = b_1^+, \quad \varphi(f_i) = \frac{[N_{i-1} + 1]}{N_{i-1} + 1} b_i^+ b_{i-1}^-, \quad i = 2, \dots, n, \quad (4c)$$

is a morphism of $U_q[sl(n+1)]$ into $W(n)$ for any $p \in \mathbf{C}$.

The proof is straightforward. In the intermediate computations the following relation is useful:

$$f(N_1, \dots, N_i, \dots, N_n) a_j^\pm = a_j^\pm f(N_1 \pm \delta_{1j}, \dots, N_i \pm \delta_{ij}, \dots, N_n \pm \delta_{nj}), \quad (5)$$

where $f(N_1, \dots, N_i, \dots, N_n) \in W(n)$ is a function of the number operators $N_1, \dots, N_i, \dots, N_n$.

We have derived the D realization (4) on the ground of an alternative to the Chevalley definition of $U_q[sl(n+1)]$ [17]. This derivation together with the expressions for (the analogs of) all Cartan-Weyl generators via CAOs will be given elsewhere.

Similar as for $sl(n+1)$, the D realization defines an infinite-dimensional representation of $U_q[sl(n+1)]$ in the Fock space $\mathcal{F}(n)$ with an orthonormed basis

$$|l\rangle \equiv |l_1, \dots, l_n\rangle = \frac{(a_1^+)^{l_1} \dots (a_n^+)^{l_n}}{\sqrt{l_1! \dots l_n!}} |0\rangle, \quad l_1, \dots, l_n = 0, 1, 2, \dots \quad (6)$$

If p is a positive integer, $p \in \mathbf{N}$, the representation is indecomposable: the subspace

$$\mathcal{F}_1(p; n) = \text{lin.env.}\{|l_1, \dots, l_n\rangle | l_1 + \dots + l_n > p\} \quad (7)$$

is an invariant subspace, whereas its orthogonal complement

$$\mathcal{F}_0(p; n) = \text{lin.env.}\{|l_1, \dots, l_n\rangle | l_1 + \dots + l_n \leq p\} \quad (8)$$

is not an invariant subspace. If $p \notin \mathbf{N}$, the representation is irreducible. In all cases however, and this is the disadvantage of the D realization, the representation of $U_q[sl(n+1)]$ in $\mathcal{F}(n)$ is not unitarizable with respect to the antilinear anti-involution $\omega : U_q[sl(n+1)] \rightarrow U_q[sl(n+1)]$, defined on the generators as

$$\omega(h_i) = h_i, \quad \omega(e_i) = f_i, \quad i = 1, \dots, n. \quad (9)$$

In order to “cure” this disadvantage we pass to introduce the HP realization.

Proposition 2 (Holstein-Primakoff realization). The linear map $\pi : U_q[sl(n+1)] \rightarrow W(n)$, defined on the generators as

$$\pi(h_1) = p - N - N_1, \quad \pi(h_i) = N_{i-1} - N_i, \quad i = 2, 3, \dots, n, \quad (10a)$$

$$\pi(e_1) = \sqrt{\frac{[N_1+1]}{N_1+1}} [p-N] a_1^-, \quad \pi(e_i) = \sqrt{\frac{[N_{i-1}]}{N_{i-1}} \frac{[N_i+1]}{N_i+1}} a_i^- a_{i-1}^+, \quad i = 2, 3, \dots, n, \quad (10b)$$

$$\pi(f_1) = \sqrt{\frac{[N_1]}{N_1}} [p-N+1] a_1^+, \quad \pi(f_i) = \sqrt{\frac{[N_{i-1}+1]}{N_{i-1}+1} \frac{[N_i]}{N_i}} a_i^+ a_{i-1}^-, \quad i = 2, 3, \dots, n, \quad (10c)$$

is a morphism of $U_q[sl(n+1)]$ into $W(n)$ for any $p \in \mathbf{C}$. If $p \in \mathbf{N}$, then $\mathcal{F}_0(p; n)$ and $\mathcal{F}_1(p; n)$ are invariant subspaces; $\mathcal{F}_0(p; n)$ carries a finite-dimensional irreducible representation; it is unitarizable with respect to the anti-involution (9) and the metric defined with the orthonormed basis (6), provided $q > 0$.

The proof is straightforward: the verification of the defining relations (2) and (3) can be carried out on a purely algebraic level. The circumstance that $\mathcal{F}(n)$ is a direct sum of its invariant subspaces $\mathcal{F}_0(p; n)$ and $\mathcal{F}_1(p; n)$ is due to the factor $\sqrt{[p - N]}$ in (10b) and $\sqrt{[p - N + 1]}$ in (10c). If $q > 0$, then $((,))$ denotes the scalar product)

$$(\pi(h_i)|l\rangle, |l'\rangle) = (|l\rangle, \pi(h_i)|l'\rangle), \quad (\pi(e_i)|l\rangle, |l'\rangle) = (|l\rangle, \pi(f_i)|l'\rangle) \quad \forall |l\rangle, |l'\rangle \in \mathcal{F}_0(p; n), \quad i = 1, \dots, n.$$

Therefore the representation of $U_q[sl(n+1)]$ in $\mathcal{F}_0(p; n)$ is unitarizable.

Let us note that the HP realization (10) of $U_q[sl(n+1)]$ can be easily expressed also in terms of deformed oscillator operators $\tilde{a}_i^\pm$, $\tilde{N}_i$, $i = 1, \dots, n$, namely operators which satisfy the relations [19-21]:

$$[\tilde{a}_i^-, \tilde{a}_j^+]_q = \delta_{ij} q^{-\tilde{N}_i}, \quad [\tilde{N}_i, \tilde{a}_j^\pm] = \pm \delta_{ij} \tilde{a}_j^\pm, \quad [\tilde{a}_i^\pm, \tilde{a}_k^\pm] = [\tilde{N}_i, \tilde{N}_k] = 0, \quad i \neq k. \quad (11)$$

From (10) and the relations between the deformed and the nondeformed operators [22]

$$\tilde{a}_i^- = \sqrt{\frac{[N_i + 1]}{N_i + 1}} a_i^-, \quad \tilde{a}_i^+ = \sqrt{\frac{[N_i]}{N_i}} a_i^-, \quad \tilde{N}_i = N_i, \quad (12)$$

one obtains a q -analogue of the HP realization for any n :

$$\pi(h_1) = p - \tilde{N} - \tilde{N}_1, \quad \pi(h_i) = \tilde{N}_{i-1} - \tilde{N}_i, \quad i = 2, 3, \dots, n, \quad (13a)$$

$$\pi(e_1) = \sqrt{[p - \tilde{N}]} \tilde{a}_1^-, \quad \pi(e_i) = \tilde{a}_i^- \tilde{a}_{i-1}^+, \quad i = 2, 3, \dots, n, \quad (13b)$$

$$\pi(f_1) = \sqrt{[p - \tilde{N} + 1]} \tilde{a}_1^+, \quad \pi(f_i) = \tilde{a}_i^+ \tilde{a}_{i-1}^-, \quad i = 2, 3, \dots, n, \quad (13c)$$

To the best of our knowledge such q -deformed analogs of HP realizations are available so far only for $U_q[sl(2)]$ [8-14] and for $U_q[sl(3)]$ [15].

Finally, adding to the generators of $U_q[sl(n+1)]$ an additional central element I , and setting $\varphi(I) = \pi(I) = p$, one obtains a D and a HP realizations of $U_q[gl(n+1)]$.

Acknowledgments

The author is thankful to Prof. S. Okubo for the kind invitation to conduct the research under the Fulbright Program in the Department of Physics and Astronomy, University of Rochester. This work was supported by the Fulbright Program, Grant No 21857.

References

- [1] Holstein T and Primakoff H 1949 *Phys. Rev.* **58** 1098
- [2] Dyson F J 1956 *Phys. Rev.* **102** 1217
- [3] Okubo S 1975 *J. Math. Phys.* **16** 528
- [4] Kittel C 1963 *Quantum Theory of Solids* (Wiley, New York)
- [5] Caspers W J 1989 *Spin Systems* (World Sci. Pub. Co., Inc., Teaneck)
- [6] Klein A and Marshalek E R 1991 *Rev. Mod. Phys.* **63** 375
- [7] Ring P and Schuck P *The Nuclear Many-Body Problem* (Springer-Verlag, New York, Heidelberg, Berlin)
- [8] Chaichian M, Ellinas D and Kulish P P 1990 *Phys. Rev. Lett.* **65** 980
- [9] Quesne C 1991 *Phys. Lett. A* **153** 303
- [10] Chakrabarti R and Jagannathan R 1991 *J. Phys. A : Math. Gen.* **24** L711
- [11] Katriel J and Solomon A I 1991 *J. Phys. A : Math. Gen.* **24** 2093
- [12] Yu Z R 1991 *J. Phys. A : Math. Gen.* **24** L1321
- [13] Kundu A and Basu Mallick B 1991 *Phys. Lett. A* **156** 175
- [14] Pan F 1991 *Chin. Phys. Lett.* **8** 56
- [15] da-Providencia J 1993 *J. Phys. A : Math. Gen.* **26** 5845
- [16] Drinfeld V 1986 *Quantum Groups (Proc. Int. Congress of Mathematics (Berkeley, 1986))*
ed A M Gleason (Providence, RI: American Physical Society) p 798
- [17] Palev T D and Parashar P 1998 *Lett.Math. Phys.* **43** 7
- [18] Polychronakos A P 1990 *Mod.Phys. Lett.* **A5** 2325
- [19] Macfarlane A J 1989 *J. Phys. A : Math. Gen.* **22** 4581
- [20] Biedenharn L C 1989 *J. Phys. A : Math. Gen.* **22** L873
- [21] Sun C P and Fu H C 1989 *J. Phys. A : Math. Gen.* **22** L983
- [22] Polychronakos A P 1990 *Mod.Phys. Lett.* **5** 2325